

Fear of Childbirth and Its Predictors in Repregnant Women After Cesarean Section: A Cross-Sectional Multicenter Study in China

Yiping Hou

Clinical Nursing Teaching and Research Section, The Second Xiangya Hospital and Xiangya School of Nursing, Central South University

Xihong Zhou (✉ zhouxihong@csu.edu.cn)

Clinical Nursing Teaching and Research Section and Department of Obstetrics, The Second Xiangya Hospital, Central South University

Min Yao

Clinical Nursing Teaching and Research Section, The Second Xiangya Hospital and Xiangya School of Nursing, Central South University

Sai Liu

Clinical Nursing Teaching and Research Section, The Second Xiangya Hospital and Xiangya School of Nursing, Central South University

Research Article

Keywords: Fear of childbirth, Predictors, Repregnant women, Cesarean section, Childbirth self-efficacy, Social support

DOI: <https://doi.org/10.21203/rs.3.rs-333746/v1>

License:  This work is licensed under a Creative Commons Attribution 4.0 International License.

[Read Full License](#)

Abstract

Background

Since the implementation of China's two-child policy in 2016, the number of repregnant women after cesarean section has increased significantly. These women are more prone to fear of childbirth compared with primiparas, which poses a great threat to their physical and mental health. However, there is currently limited research on the problem in China. The aim of this study was to assess fear of childbirth and its predictors in repregnant women after cesarean section in China.

Methods

A cross-sectional multicenter study was conducted in three hospitals from June 7 to December 7, 2020, in Changsha, China. A convenience sampling technique was employed to include participants. Pretested and organized questionnaires were used to collect data. After checking for completeness, data were exported to statistical software for analysis. Both univariate analysis and multiple linear regression analysis were computed to assess fear of childbirth and its predictors. Statistical significance was declared at a P-value of < 0.05 .

Results

A total of 358 repregnant women after cesarean section have participated in this study. The average score of fear of childbirth was 43.76 (standard deviation = 5.27, range 17–58). Number of cesarean sections, experience with previous cesarean section, childbirth self-efficacy and social support were significantly associated with fear of childbirth.

Conclusions

In this study, repregnant women after cesarean section in China had moderate fear of childbirth, and the number of cesarean sections, experience with previous cesarean section, childbirth self-efficacy and social support were predictors of fear of childbirth. It is important for healthcare professionals to find repregnant women after cesarean section at high risk of fear of childbirth and provide appropriate services during pregnancy.

Background

Fear of childbirth (FOC) is described as a negative feeling before, during or after delivery when thinking about future delivery or experiencing others' fearful responses to childbirth and labor pain [1]. In other words, FOC is a negative cognitive assessment of the pathological dread and avoidance of childbirth [2] and manifests in the form of physical discomfort, nightmares, and difficulty concentrating on work and

family activities [3]. In addition, FOC may develop during pregnancy, and the content of such fear may include pain, uncertainty, loss of control, and the possibility of having an impaired or stillborn child [4]. It is not uncommon for pregnant women to fear childbirth, as it has been estimated that FOC affects approximately 14% of pregnant women worldwide [5]. Studies in China have also shown that the incidence of FOC among women is 50% or higher [6]. The degree of FOC varies from person to person, ranging from mild to severe fear. Research has reported that 20% of pregnant women experience moderate FOC and that 6–13% of pregnant women experience severe FOC [7].

FOC has many negative effects on women's physical and mental health, including pregnancy complications, severe pain and the use of anesthesia during childbirth, prolonged childbirth, mother-child relationship difficulties, postpartum depression and posttraumatic stress disorder [7–13]. In addition, FOC is often the reason why women request a cesarean section (CS), which may lead to the occurrence of a CS without medical indications [4], resulting in a waste of medical resources and increasing socioeconomic burdens. Increasing evidence has shown that FOC is related to the mode of delivery [14, 15]. A large cohort study of more than 700,000 pregnant women found that the CS rate of women with FOC was 4.4 times higher than that of women without FOC [12]. Studies have indicated that young, low-educated women are more likely to suffer from FOC compared to older, high-educated women [16]. In China, high obstetric intervention rates and the lack of high-quality maternal care may cause pregnant women to fear childbirth. Fear of pain and a previous difficult delivery have also been found to be related to FOC [17].

Since the implementation of China's two-child policy in 2016, the number of pregnant women has increased significantly [18]. The high rate of CS (40–60%) in the past has increased the number of repregnant women after CS [19]. These women are a special group compared to primiparas due to their history of scarred uterus leading to a more complicated delivery process, which makes them more incline to FOC [20]. Moreover, these women not only have the same FOC as that of primiparas but also the fear that their previous CS may cause various complications in their current pregnancy and childbirth: postpartum hemorrhage, infection, placenta previa, and placenta accreta [21].

Against the special background of the two-child policy and high CS rates in China, it is important for healthcare providers to understand, recognize and address FOC in repregnant women after CS. Although this problem requires urgent attention, to the best of our knowledge, in China, (1) most studies have focused on anxiety and depression, while the research on FOC is limited; (2) studies have concentrated on primiparas and have often ignored repregnant women after CS; and (3) furthermore, the available studies on FOC have been conducted in high-income areas, and thus, little is known about this problem in low-income areas. Given the limited understanding of FOC in repregnant women after CS in China, this study aimed to assess FOC and its predictors.

Methods

Study design and setting

A cross-sectional descriptive study was conducted from June 7 to December 7, 2020, in Changsha, China. Changsha is a subprovincial city located in the south-central part of China, and is the capital of Hunan Province, and has a population of approximately 8 million. The birth rate in Changsha is over 5,000 babies per year.

Participants

Women during the third trimester of pregnancy who were receiving antenatal care at the study hospitals during the study period were recruited. The inclusion criteria were as follows: (1) being older than 18 years, (2) having a history of CS(s); (3) being able to understand the content of the questionnaires, and (4) providing consent to participate. Women with concurrent major physical or mental health problems were excluded.

Sample size determination and sampling procedure

The required sample size was determined using a single population formula based on the assumption of a 95% confidence interval with a margin of error of 5%. Three hospitals were chosen (The Second Xiangya Hospital, Hunan Provincial People's Hospital and Changsha Maternal and Child Health Hospital) that are purposively dependent on antenatal services and obstetrics care. The annual delivery rate of each hospital is approximately 4,000 live births. The sample was apportioned to every hospital by investigating the number of deliveries earlier in the year until the complete sample was obtained.

Materials

In this study, the Childbirth Attitude Questionnaire (CAQ), which was developed by Lowe in 2000 [22], was used to measure FOC. The CAQ is a 16-item questionnaire that uses a 4-point Likert scale. The item scores are summed to provide a total score (range 16-64), with higher scores indicating higher levels of FOC. The Chinese version of the CAQ has been used with good reliability and validity [23]. The Cronbach's alpha for the scale was 0.92. For the classification of no, low, moderate and severe levels of FOC, the cutoff values were 16, 28, 40, and 52, respectively.

The short form of the 32-item Chinese Childbirth Self-Efficacy Inventory (CBSEI-C32) [24] was used in this study to measure childbirth self-efficacy. The scale has two parallel subscales—the outcome expectancy subscale and the efficacy expectancy subscale—which consist of the same 16 items measuring coping behavior for childbirth on a 4-point Likert scale. The sum of the each subscale is the total score (range 32-128), with higher scores indicating higher levels of childbirth self-efficacy. The scale has a reported Cronbach's alpha value of 0.90 and has been shown to be a valid and reliable measure in Mainland China [24].

The Social Support Rating Scale (SSRS), which was developed by Xiao [25], was used to measure social support in this study. There are 10 items in this scale, including three dimensions—objective support (3 items), subjective support (4 items) and the utilization of social support (3 items). The sum of each

dimension is the total score, with higher scores indicating higher levels of social support. The scale has Cronbach's alpha values of 0.825-0.896 and has been used widely in China [25].

Demographic and obstetric data included age (<35/≥35), pregnant women who were the only child in their families (no/yes), residential area (rural/urban), education (elementary school and below/junior high school/senior high school/college and above), occupation (office clerk/agricultural worker/self-employed/freelance), self-rated economic status (poor/fair/good), number of CSs (1/≥2), and experience with previous CS (no or mild fear/moderate or severe fear).

Procedures and ethical consideration

Ethical approval was obtained from the university and the study hospital. Participation in the study was voluntary, and all participants provided written informed consent before participation.

After obtaining ethical approval, a pilot study was conducted with 10 eligible subjects to evaluate the feasibility of the study and identify any unpredictable data collection problems, and no problems were reported. Training for data collectors was provided to ensure the accuracy and consistency of data collection. All eligible subjects who were waiting for their appointment in the obstetric clinic of the research hospital were invited to participate in the study. After providing informed consent, participants were asked to complete the demographic and obstetric questionnaire, the CAQ, the CBSEI-C32 and the SSRS. Data collectors remained in the vicinity to answer questions and personally collected the returned questionnaires.

Data processing and analysis

After data collection, all collected questionnaires were checked for completeness and internal consistency to exclude missing or inconsistent data. Data were entered and analyzed using IBM SPSS for Windows, version 16.0 (SPSS Inc., Chicago, IL, USA). Mean (M), standard deviation (SD), frequency and percentage were used to describe demographic and obstetric data, CAQ, CBSEI-C32 and SSRS scores. Independent sample *t*-test and analysis of variance were used to compare the CAQ scores with different demographic and obstetric characteristics. Pearson correlation coefficients were calculated to measure the relationships among CBSEI-C32, SSRS, and CAQ. If $p < 0.05$ in the above tests, then significant variables were entered in the multiple linear regression analysis to predict FOC. The variance inflation factor (VIF) was used to assess multicollinearity among the predictors.

Results

Sample and demographic characteristics of participants

A total of 410 eligible women visited the clinics during the period of investigation. Forty-one women (10%) refused to participate and 11 women (2.68%) consented but did not complete the questionnaires, resulting in a total of 358 pregnant women being included in this study. The ages of these women ranged

from 22 to 44 years ($M=33.60$, $SD= 4.02$), and their M gestational age was 253.82 days ($SD=17.93$) at enrollment. All the participants were married.

CAQ scores

The average CAQ score among these repregnant women was 43.76 ($SD=5.27$, range 17-58). The M score of each item of the CAQ was 2.73 ($SD=0.55$). The women reported their greatest fears as being the baby being injured during the childbirth and having painful labor contractions. The least fears that women reported were having nightmares about childbirth and the hospital environment (Table 1).

Demographic and obstetric characteristics and their association with the CAQ

Of the pregnant women studied, 72.91% ($n=261$) were younger than 35 years, and 32.96% ($n=118$) were only child in their families. The majority (63.69%, $n=228$) of the women lived in urban areas, and 14.81% ($n=53$) had an educational level of college or above. Approximately 36.03% of respondents ($n=129$) were office clerks, and 49.16% ($n=176$) reported good economic status. The minority (8.66%, $n= 31$) of these women had more than 2 CS events, and 33.8% ($n=77$) had moderate or severe fear of their previous CS. Pregnant women who were the only child in their families, self-rated economic status, number of CSs, and experience with previous CS were significantly associated with the CAQ (Table 2).

Correlations among the CBSEI-C32, SSRS, and CAQ

The average total objective value of the CBSEI-C32 among respondents was 87.59 ($SD=18.81$, range 32-128). The average total SSRS score among these women was 45.35 ($SD=7.97$, range 17-60). Pearson correlation coefficients indicated that CBSEI-C32 and SSRS were significantly negatively related to the CAQ, and CBSEI-C32 was significantly positively related to SSRS. (Table 3).

Predictors of FOC

Based on the significant results between candidate predictors and the CAQ, these variables (pregnant women who were the only child in their families, self-rated economic status, number of CSs, experience with previous CS, CBSEI-C32 and SSRS) were validated through multivariate linear regression (Table 4), which showed that number of CSs, experience with previous CS, childbirth self-efficacy and social support were predictors of FOC. The VIF value was less than 10, which indicated that there was no multicollinearity among the variables. These variables accounted for 83.1% of the total variance. The analysis was found to be statistically significant ($F = 440.77$, $p < 0.000$) (Table 5).

Discussion

We used the CAQ to assess the FOC in repregnant women after CS in China. The total CAQ score in our study was 43.76, which was close to that in a study conducted in Turkey [26] and higher than those in studies conducted in the U.S. (31.70) and Greece (31.22) [22, 27]. This result indicated that repregnant women after CS in China might have the same moderate FOC as do women in Turkey, which was higher

than that of pregnant women in the U.S. and Greece. This finding may be related to different study populations. Repregnant women after CS in our study were more likely to have FOC due to their history of CS, leading to complicated childbirth compared with primiparas in the U.S. and Greek studies.

The M score of each item was 2.73, which was higher than that in a study conducted in Thailand (2.39) [28]. This finding may be related to the implementation of the two-child policy leading to more repregnant women after CS being older in age (≥ 35 years) in our study. Relevant studies have suggested that older age is associated with FOC [12, 29]. In our study, 27.1% of women were older, whereas in the abovementioned study in Thailand, only 24.3% of women were older. Furthermore, our study indicated that repregnant women after CS had the most fears for their baby's health and painful labor contractions, a finding that was consistent with a study by Lowe et al. [22]. Therefore, the CAQ can be used to identify the specific areas of pregnant women's FOC. Moreover, appropriate intervention can target the areas where pregnant women obtained high scores.

Our research found that the number of CSs (≥ 2) was positively correlated with FOC. Until now, no study presenting such a finding has been published in English. A possible reason for this may be that most studies were focused on primiparas' FOC, whereas our study included a sample of repregnant women with a history of CS(s). Against the specific background of the two-child policy and high CS rates in China [18, 19], the significance of this risk factor cannot be ignored. A related study found that the degree of FOC in repregnant women after CS was higher than that of primiparous women [20]. A CS is a traumatic surgical procedure, and women thus bear potential risks. Experience with a previous CS affects women's future reproductive ability and increase their risk of ectopic pregnancy and placental implantation [30]. Moreover, repeated CSs (≥ 2) may cause uterine rupture in the perinatal period because of a scarred uterus [31]. These hazards greatly increase FOC in repregnant women after CS. Thus, clinicians should detect repregnant women with a history of repeated CSs at high risk for FOC and initiate or refer them to appropriate services during pregnancy.

This study showed that the experience of women with previous CS (moderate or severe fear) was positively related to FOC, which was in alignment with the results of studies conducted in Turkey and Hungary [32, 33]. Repregnant women after CS with negative delivery experience were prone to FOC. The reason may be that women who had negative childbirth experience in a previous CS were worried that a similar negative experience may occur in their next childbirth experience. Fenwick et al. reported that women who experienced problems such as miscarriage or preeclampsia in their previous pregnancy were most afraid of facing similar or more severe conditions this time around [34]. A qualitative study demonstrated that an inadequate understanding and incorrect information about delivery can lead to fear among women [35]. Melender et al. found that negative stories and experiences shared by others were also one of the sources of FOC [13]. In addition, women were exposed to more negative information, which may have had negative effects on their thoughts about childbirth and may have ultimately increased their FOC. Previous studies showed that the occurrence of situations such as a negative birth experience and birth trauma could lead to FOC among women [36, 37]. However, Phunyammalee et al. found that a similar FOC between women with and without previous CS [38], which may be attributable to

the fact that women received better care in their previous CSs, resulting in a less negative experience. Negative childbirth experiences can cause women to hold negative emotions about future childbirth, which can affect their choice of delivery mode [39]. In contrast, positive childbirth experiences can enhance self-esteem, satisfaction and the desire to be pregnant again [40]. Therefore, it is clear that women with negative delivery experience should receive more support from professionals during pregnancy.

Our study indicated that childbirth self-efficacy was negatively related to FOC, which was consistent with self-efficacy theory [41]. Bandura defined self-efficacy as a dynamic cognitive process in which individuals evaluate their ability to cope with different realities and execute required behaviors [42]. Moreover, previous studies conducted by Salomonsson et al. and Tanglakmankhong et al. also found that childbirth self-efficacy was related to FOC [28, 43]. According to self-efficacy theory, emotional arousal is one of the sources of self-efficacy [41]. Repregnant women after CS tend to have negative emotions during childbirth, which may be related to their concerns about labor pains, uterine rupture, and fetal health. The arousal of disgust caused by threatening situations usually reduces self-efficacy. Women with low self-efficacy may have a limited ability to generate motivation to cope with delivery. These women treat childbirth as a difficult task and wonder if they are capable of coping with it. In contrast, women with higher childbirth self-efficacy reported lower levels of FOC [22]. In addition, previous studies have reported that self-efficacy is a determinant of FOC in pregnant women [43], indicating that improvement of the self-efficacy of pregnant women can increase their confidence during childbirth and may help reduce their FOC.

In addition to the abovementioned factors, we also observed that social support was negatively correlated with FOC a finding that was similar to those of other studies [44, 45]. Social support is a kind of interpersonal relationship, which refers to the provision of material and spiritual help and supportive behaviors by family members, neighbors, colleagues, relatives, and the treatment team toward individuals [46]. Relatives, especially spouses, show their respect for pregnant women by supporting them, thus increasing their hope. A related study showed that a lack of social support or expressed dissatisfaction with one's partner was also predictive of FOC. Moreover, the more dissatisfied women were with their partnership and the lack of social support, the more fearful they were of childbirth [3]. Social support is an important factor in maintaining individual mental health during pregnancy. The study by Fisher et al. showed that social relationships and the strong support of midwives can enhance women's beliefs that childbirth is a physiological and controllable process, thereby improving their mental health and reducing their FOC [47]. Adequate social support helps pregnant women acquire pregnancy-related knowledge [38]. However, pregnant women with limited social support may have limited access and ability to obtain relevant and correct information, whether provided by clinicians or found online information. Insufficient information about the delivery process can lead to increased FOC. Therefore, actual information about the need for more social support during pregnancy is crucial for helping pregnant women actively relieve their FOC and approach delivery.

Limitations

This study has several limitations that must be noted. First, this is a cross-sectional study, which limits the possibility of establishing causal relationships between variables. Longitudinal and experimental studies are warranted to determine the relevant factors of FOC in repregnant women after CS in China. Second, this study was conducted in three university hospitals in a particular city, so it cannot be representative of the majority of pregnant women in China. Further research is needed on women in rural and underserved areas of the country. In addition, effective interventions for reducing FOC should be developed and evaluated in the future.

Conclusions

In this study, repregnant women after CS in China had moderate FOC, and the number of CSs, experience with previous CS, childbirth self-efficacy and social support were predictors of FOC. The identification of the predictors of FOC in repregnant women after CS in China assists in the identification of those who may have FOC. The early identification of women with FOC will allow healthcare professionals to provide appropriate interventions to reduce FOC during pregnancy, which will decrease the risk of negative psychological and obstetric consequences in these women in the future.

List Of Abbreviations

FOC

Fear of childbirth; CS:Cesarean section; M:Mean; SD:Standard deviation; CAQ:Childbirth Attitude Questionnaire; CBSEI-C32:32-item Chinese Childbirth Self-Efficacy Inventory; SSRS:Social Support Rating Scale; VIF:Variance inflation factor; B:Partial regression coefficient; SE:Standard error.

Declarations

Ethics approval and consent to participate

The study was approved by the Ethics Review Committee of Nursing and Behavioral Medicine Research, Xiangya School of Nursing, Central South University (no. E201915). This study was conducted in accordance with the guidelines of the Declaration of Helsinki. Participation in the study was voluntary, and all participants provided written informed consent before participation.

Consent for publication

Not applicable.

Availability of data and materials

The datasets used and/or analyzed during the current study are available from the corresponding author on reasonable request.

Competing interests

The authors declare that they have no competing interests.

Funding

This work was supported by the Fundamental Research Funds for the Central Universities of Central South University [grant number 2019zzts193]. The funding body played no role in study design, data collection, analysis or interpretation and writing of the manuscript.

Authors' contributions

All the authors contributed to the work and approved the final version of this manuscript. This study was designed by YPH, XHZ, MY and SL. SL and MY performed the statistical analysis. YPH wrote the manuscript. XHZ reviewed and edited manuscript. All authors read and approved the manuscript.

Acknowledgements

We would like to thank The Second Xiangya Hospital, Xiangya School of Nursing, Central South University, Hunan Provincial People's Hospital and Changsha Maternal and Child Health Hospital for letting us to conduct this study. Our deepest gratitude also goes to study participants, data collectors, and supervisor for their invaluable support to make this study real.

References

1. Nilsson C, Hessman E, Sjöblom H, et al. Definitions, measurements and prevalence of fear of childbirth: a systematic review. *BMC Pregnancy Childbirth*. 2018;18(1): 28.
2. Hofberg, K., Brockington, I.. Tokophobia: an unreasoning dread of childbirth. A series of 26 cases. *Br. J. Psychiatry*. 2000;176,83–85.
3. Saisto T, Salmela-Aro K, Nurmi JE, et al. Psychosocial characteristics of women and their partners fearing vaginal childbirth. *BJOG*. 2001;108(5):492–498.
4. Rouhe H, Salmela-Aro K, Toivanen R, et al. Obstetric outcome after intervention for severe fear of childbirth in nulliparous women-randomised trial. *BJOG*. 2013; 120(1):75–84.
5. O'Connell MA, Leahy-Warren P, Khashan AS, et al. Worldwide prevalence of tocophobia in pregnant women: systematic review and meta-analysis. *Acta Obstet Gynecol Scand*. 2017;96(8):907–920.
6. Jia Z, Yang YY, Qiu Y. Analysis on the childbirth attitudes of primiparous women and related influencing factors. *Nursing of Integrated Traditional Chinese and Western Medicine*. 2019;5(6):67–70.
7. Rouhe, H., Salmela-Aro, K., Halmesmäki, E.. et al. Fear of childbirth according to parity, gestational age, and obstetric history. *BJOG: Int. J. Obstet. Gynaecol*. 2009; 116(1),67–73.

8. Adams S, Eberhard-Gran M, Eskild A. Fear of childbirth and duration of labour: a study of 2206 women with intended vaginal delivery. *BJOG: An International Journal of Obstetrics Gynaecology*. 2012;119(10):1238–1246.
9. Vanden Bussche E, Crombez G, Eccleston C, et al. Why women prefer epidural analgesia during childbirth: the role of beliefs about epidural analgesia and pain catastrophizing. *European journal of pain*. 2007;11(3):275–282.
10. Alehagen S, Wijma B, Wijma K. Fear of childbirth before, during, and after childbirth. *Acta obstetrica et gynecologica Scandinavica*. 2006;85(1):56–62.
11. Mazúchová L, Škodová Z, Kelčíková S, et al. Factors Associated With Childbirth-Related Fear Among Slovak Women. *Central European Journal of Nursing Midwifery*. 2017;8(4):742–748.
12. Raisanen S, Lehto SM, Nielsen HS, et al. Fear of childbirth in nulliparous and multiparous women: a population-based analysis of all singleton births in Finland in 1997–2010. *BJOG*. 2014;121(8):965–970.
13. Melender HL. Experiences of fears associated with pregnancy and childbirth: a study of 329 pregnant women. *Birth*. 2002;29(2):101–111.
14. Ryding EL, Lukasse M, Parys AS, et al. Fear of childbirth and risk of cesarean delivery: a cohort study in six European countries. *Birth*. 2015;42(1):48–55.
15. Stoll K, Edmonds JK, Hall WA. Fear of childbirth and preference for cesarean delivery among young American women before childbirth: a survey study. *Birth*. 2015;42(3):270–276.
16. Laursen, M., Hedegaard, M., Johansen, C.. Fear of childbirth: predictors and temporal changes among nulliparous women in the Danish national birth cohort. *BJOG: Int. J. Obstet. Gynecol*. 2008;115(3):354–360.
17. El-Aziz SNA, Mansour SE-S, Hassan NF. Factors associated with fear of childbirth: It's effect on women's preference for elective cesarean section. *Journal of Nursing Education and Practice*. 2017;7(1):133–146.
18. Meng X, Lin P. Differences in population distribution between high-risk pregnant women and non-open child policy. *Matern Child Health Care China*. 2016.;31(20):4266–4268.
19. Liu XJ, Jia GF, Wen Y. Analysis of 10-year cesarean section rate and indication of cesarean section. *Matern Child Health Care China*. 2015;30(4):551–553.
20. SLUIJS A M, CLEIREN M, SCHERJON S A, et al. No relationship between fear of childbirth and pregnancy-delivery-outcome in a low-risk Dutch pregnancy cohort delivering at home or in hospital. *J Psychosom Obstetr Gynecol*. 2012;33(3):99–105.
21. Yang KY. Psychological characteristics and social factors of women choosing vaginal birth after cesarean. D. Xinjiang: Shihezi University. 2017:1–43.
22. Lowe NK. Self-efficacy for labor and childbirth fears in nulliparous pregnant women. *J Psychosom Obstetr Gynaecol*. 2000;21(4):219–224.

23. Wei J, Liu F, Zhang LF, et al. Reliability and validity test of the Chinese version of the Childbirth Attitude Questionnaire. *J Nurs*. 2016;31(2):81–83.
24. Gao LL, Ip W Y, Sun K. Validation of the short form of Chinese Childbirth Self- efficacy Inventory in mainland China. *Res Nurs Health*. 2011;34,49–59.
25. Xiao S, Yang D, Hui S, et al. Effects of social support on physical and psychological health. *Chin Ment Health J*. 1987;4(1):183-187.
26. D Coşkuner Potur, R Mamuk, N H Şahin, et al. Association between fear of childbirth and maternal acceptance of pregnancy. *Int Nurs Rev*. 2017;64(4): 576–583.
27. Gourounti K, Kouklaki E, Lykeridou K. Validation of the Childbirth Attitudes Questionnaire in Greek and psychosocial characteristics of pregnant women with fear of childbirth. *Women Birth*. 2015;28(3):44–51.
28. Tanglakmankhong, K., Perrin, N.A., Lowe, N.K.. Childbirth Self-Efficacy Inventory and Childbirth Attitudes Questionnaire: psychometric properties of Thai Language versions. *J. Adv. Nurs*. 2011;67(1),193–203.
29. Gao GF. Fear of childbirth and its influencing factors in primiparous women of childbearing age. *Maternal and Child Health Care of China*. 2020;35(3):541 – 43.
30. Xie W. Discussion on related problems of vaginal delivery and cesarean section. *Journal of Yanan University (Med Sci)*. 2012;10(4):27–28.
31. Hou YP, Zhou XH, Shi LH. Influence factors and pregnancy outcomes for pernicious placenta previa with placenta accreta. *J Cent South Univ (Med Sci)*. 2020;45(9):1074–1081.
32. Pınar Serçekuş, Okan Vardar, Sevgi Özkan. Fear of childbirth among pregnant women and their partners in Turkey. *Sexual & Reproductive Healthcare*. 2020;24: 1–6.
33. Vahideh MoghaddamHosseini, Alexandra Makai, Katalin Varga. Assessing fear of childbirth and its predictors among Hungarian pregnant women using Wijma Delivery Expectancy/Experience Questionnaire subscales. *Psychology, Health & Medicine*. 2019;24(7):879–889.
34. Fenwick J, Bayes S, Johansson M. A qualitative investigation into the pregnancy experiences and childbirth expectations of Australian fathers-to-be. *Sex Reprod Healthc*. 2012;3(1):3–9.
35. Serçekus P, Okumus H. Fears associated with childbirth among nulliparous women in Turkey. *Midwifery*. 2009;25(2):155 – 62.
36. Dencker A, Nilsson C, Begley C, Jangsten E, Mollberg M, Patel H, et al. Causes and outcomes in studies of fear of childbirth: a systematic review. *Women Birth*. 2019;32(2):99–111.
37. Nilsson C, Bondas T, Lundgren I. Previous birth experience in women with intense fear of childbirth. *J Obstet Gynecol Neonatal Nurs*. 2010;39(3):298–309.
38. Phunyammalee M, Buayaem T, Boriboonhirunsarn D. *J Obstet Gynaecol*. 2019;39(6):763–767.
39. Takegata M, Haruna M, Matsuzaki M, et al. Translation and validation of the Japanese version of the Wijma Delivery Expectancy/Experience Questionnaire version A. *Nurs Health Sci*. 2013,15(3):326–332.

40. O'Hara M W,McCabe J E. Postpartum depression: current status and future directions. *Annu Rev Cli Psychol.* 2013;9:379–407.
41. Bandura, A.. Self-efficacy: toward a unifying theory of behavioral change. *Psychol. Rev.* 1977;84(2),191–215.
42. A. B. Self-efficacymechnism in human agency. *Am Psychol.* 1982;37 (2):122 – 47.
43. Salomonsson, B., Gullberg, M.T., Alehagen, S., et al. Self-efficacy beliefs and fear of childbirth in nulliparous women. *J. Psychosom. Obstet. Gynaecol.* 2013;34(3),116 – 21.
44. Lukasse M, Schei B, Ryding EL,et al. Prevalence and associated factors of fear of childbirth in six European countries. *Sexual & Reproductive Healthcare.* 2014;5(3):99–106.
45. Toohill J, Creedy DK, Gamble J,et al. A cross-sectional study to determine utility of childbirth fear screening in maternity practice-an Australian perspective. *Women and Birth.* 2015;28(4):310–316.
46. Nazare S, Salare P, Mazlom R, Mehram F. The comparative study of social support postpartum mothers multiparity in the first and second month after delivery. *Public Health Research Khorasa.* 2014;8:1–9.
47. Fisher C, Hauckb Y, Fenwickb J. How social context impacts on women's fears of childbirth. *Soc Sci Med.* 2006;63(1):64–75.

Tables

Table 1
CAQ scores listed by item ($n = 358$)

Original item no.	CAQ items	M (SD)
11	I have a fear of my baby being injured during childbirth.	3.42 (0.79)
12	I have a fear of painful labor contractions.	3.41 (0.78)
9	I have a fear of needing a cesarean section.	3.40 (0.79)
6	I have some fear of something being wrong with my baby.	3.31 (0.88)
2	I am truly afraid of giving birth.	3.19 (0.76)
16	Overall, I would rate my anxiety about childbirth as 1 (no anxiety), 2 (low anxiety), 3 (moderate anxiety), or 4 (high anxiety).	3.16 (0.77)
10	I have a fear of being torn during the birth of my baby.	2.95 (0.82)
4	I have a fear of bleeding too much during childbirth.	2.89 (0.84)
1	I have a fear of losing control of myself during childbirth.	2.71 (0.89)
13	I have difficulty relaxing when thinking of the upcoming birth.	2.54 (0.66)
7	I have a fear of painful injections.	2.33 (0.57)
8	I have a fear of being left alone during labor.	2.22 (0.71)
5	I have a fear that I will not be able to help during childbirth.	2.16 (0.72)
15	I have a fear of not getting the kind of care that I want.	2.13 (0.81)
14	I have a fear of the hospital environment.	2.08 (0.58)
3	I have nightmares about childbirth.	1.85 (0.65)

Abbreviations: CAQ: Childbirth Attitude Questionnaire; M: mean; and SD: standard deviation.

Table 2
Demographic and obstetric characteristic and comparisons of CAQ scores among subgroups (*n* = 358)

Variables	Total n (%)	CAQ M (SD)	<i>F</i> or <i>t</i>	<i>P</i>
Age (years)			-1.81	0.071
< 35	261 (72.91)	43.46 (5.30)		
≥ 35	97 (27.09)	44.59 (5.13)		
The only child in her family			3.42	0.001
No	240 (67.04)	43.10 (5.36)		
Yes	118 (32.96)	45.10 (4.85)		
Residential area			0.13	0.894
Rural	130 (36.31)	43.81 (4.18)		
Urban	228 (63.69)	43.74 (5.82)		
Education			1.82	0.143
Elementary school and below	39 (10.89)	42.69 (4.87)		
Junior high school	60 (16.76)	43.63 (7.04)		
Senior high school	206 (57.54)	43.65 (4.80)		
College and above	53 (14.81)	45.15 (4.85)		
Occupation			1.69	0.168
Office clerk	129 (36.03)	44.12 (4.58)		
Agricultural worker	13 (3.63)	43.00 (4.60)		
Self-employed	89 (24.86)	42.74 (6.46)		
Freelance	127 (35.48)	44.19 (5.01)		
Self-rated economic status			4.16	0.006
Poor	41 (11.45)	46.34 (5.08)		
Fair	141 (39.39)	43.78 (5.13)		
Good	176 (49.16)	43.15 (5.27)		
Number of CSs			-3.60	0.000

Abbreviations: CAQ: Childbirth Attitude Questionnaire; M: mean; SD: standard deviation; and CS: cesarean section.

Variables	Total n (%)	CAQ M (SD)	For t	P
1	327 (91.34)	43.46 (5.15)		
≥ 2	31 (8.66)	46.97 (5.54)		
Experience with previous CS			-3.37	0.001
No or mild fear	281 (66.20)	43.28 (5.36)		
Moderate or severe fear	77 (33.80)	45.53 (4.55)		
Abbreviations: CAQ: Childbirth Attitude Questionnaire; M: mean; SD: standard deviation; and CS: cesarean section.				

Table 3
Relationships among the CBSEI-C32, the SSRS, and CAQ scores ($n = 358$)

Variable	CAQ	CBSEI-C32	SSRS
CAQ	1	-0.905**	-0.869**
CBSEI-C32	-0.905**	1	0.979**
SSRS	-0.869**	0.979**	1
Abbreviations: CAQ: Childbirth Attitude Questionnaire; CBSEI-C32: 32-item Chinese Childbirth Self-Efficacy Inventory; and SSRS: Social Support Rating Scale. ** $p < 0.01$.			

Table 4
Independent variable assignment of factors associated with FOC

Variable	Assignment
The only child in her family	1 = no; 2 = yes
Self-rated economic status	1 = poor; 2 = fair; 3 = good
Number of CSs	1 = 1; 2 = ≥ 2
Experience of previous CS	1 = no or mild fear; 2 = moderate or severe fear
CBSEI-C32	Original value
SSRS	Original value
Abbreviations: FOC: fear of childbirth; CS: cesarean section; CBSEI-C32: 32-item Chinese Childbirth Self-Efficacy Inventory; and SSRS: Social Support Rating Scale.	

Table 5
Multiple linear regression analysis of predictors of FOC.

Variable	<i>B</i>	<i>SE</i>	<i>t</i>	<i>p</i>	VIF
Constant	66.905	1.137	58.866	0.000	-
CBSEI-C32	-0.360	0.030	-12.092	0.000	3.86
Number of CSs	1.333	0.449	2.969	0.003	1.22
SSRS	-0.234	0.071	-3.312	0.001	4.01
Experience with previous CS	1.623	0.302	2.061	0.040	1.18
<i>F</i> = 440.77, <i>p</i> < 0.000; <i>R</i> ² = 0.833, <i>R</i> _{adj} = 0.831.					
Abbreviations: <i>B</i> : partial regression coefficient; <i>SE</i> : standard error; FOC: fear of childbirth; CBSEI-C32: 32-item Chinese Childbirth Self-Efficacy Inventory; CS: cesarean section; SSRS: Social Support Rating Scale; and VIF: variance inflation factor.					